

Guide to Terrestrial Noxious and Invasive Weeds

Near Lake Tulloch Reservoir



Public Identification, Prevention & Management Guide

Terrestrial Noxious Weeds Near Tulloch Reservoir - 2026



Purpose

The purpose of this Noxious Weed pamphlet is to assist the Tri-Dam Project in reaching the public to enlist their help in controlling the invasive, alien weeds that are threatening to displace native plants in the natural environment at Tulloch Reservoir. These invasive alien weeds also threaten to displace rare plant populations and reduce wildlife habitat values.

Tri-Dam's FERC License Conditions

The Federal Energy Regulatory Commission (FERC) required specific conditions to be implemented for the renewal of the FERC License for Tri-Dam's Tulloch Hydroelectric Project. FERC required a Vegetation Management Plan to include a plan to manage noxious weeds, in FERC's Article 405, at pages 23- 24, of the Order Issuing New License on February 16, 2006, for the South San Joaquin Irrigation and Oakdale Irrigation Districts' Tri-Dam Project No. 2067-020. The Vegetation Management Plan was approved on July 1, 2008. As part of the plan, a terrestrial noxious weed survey is performed annually by Tri-Dam Staff. Once identified, staff perform mitigation efforts where feasible. To help Tri-Dam prevent the spread of noxious weeds, this pamphlet informs the public around Tulloch Reservoir.

The following plants were found at Tulloch Reservoir by Project biologists and are classified by state and federal agencies as invasive or "Noxious Weeds". These weeds occur in patches on federal lands, state lands, TriDam lands, and private lands around the Tri-Dam Project boundary at Tulloch Reservoir:

1. Barbed Goatgrass (*Aegilops triuncialis*)
2. Chinese Tree-of-Heaven (*Ailanthus altissima*)
3. Italian Thistle (*Carduus pycnocephalus*)
4. Slender-flower Thistle (*Carduus tenuiflorus*)
5. Tocalote [Malta Starthistle] (*Centaurea melitensis*)
6. Yellow Star Thistle (*Centaurea solstitialis*)
7. Bull Thistle (*Cirsium vulgare*)
8. Klamath Weed [St. John's Wort] (*Hypericum perforatum*)
9. Himalayan Blackberry (*Rubus discolor* [procerus] [armeniacus])
10. Medusa-head Grass (*Taeniantherum* [Elymus] caput-medusae)

Noxious Weed Control Methods

The following pages discuss the biology of the plants listed by State or Federal agencies as "Noxious Weeds" and the control methods that have been reported as successful for each plant. The different methods for controlling weeds are summarized in this pamphlet.

Weed Prevention

The first line of defense against noxious weed invasions is to prevent their introduction into the project area. The following preventive measures should be utilized:

- Avoid disturbance to natural areas, including clearing of native vegetation and planting of nonnative plants.
- Avoid entering areas with existing noxious weed populations, or if entry is necessary, conduct work first in clean areas and then in areas with weeds, to avoid spreading weeds within the project area.
- Remember to clean equipment before leaving infested areas. Do not dump yard waste containing weeds that can spread seeds or re-root. Dispose of weed seed heads in the garbage. Do not mulch invasive weeds.
- Do not purchase or use invasive exotic species in landscaping or for land restoration or erosion control projects.
- Use hay for erosion control that is not contaminated with weeds. Commit to using certified weed-free straw for all construction or restoration projects or, if unavailable, use rice straw.
- For landscaping, use plants that are native to the Central Sierra foothills region as much as possible or those that are not known to be invasive.
- When planting seeds, use weed-free seed to help prevent the spread of most noxious weeds.
- Know your plants. If you are unsure of the identification of a plant, take a sample to a university, arboretum, department of agriculture office, local nature center, or native plant society for assistance. Find out whether it is known or suspected to be invasive. If the exotic plant is closely related to an invasive species, it is likely to have similar tendencies. To be on the safe side, if you don't know it, don't grow it.
- Avoid buying nursery stock that appears unhealthy and discard any unhealthy nursery stock in the trash. Do not mulch these plants on the project site.
- Control exotic invasive plants in your landscape either by removing them entirely or by managing them to prevent their spread outside your property. This may include pruning to prevent flowering and seed dispersal or cutting, mowing, or herbicide use to prevent vegetative spread.
- Do not dispose of aquatic pond or aquarium plants in natural waterways or lakes.
- Clean off your boat's propeller and boat trailer of any plant material before placing it in the lake after boating in other waterways.
- Discuss your concerns about invasive exotic plants with nurseries and garden shops and ask them not to sell these species. Provide them with printed material (such as this) explaining the problem to read later. Ask for non-invasive alternatives instead.

Noxious Weed Control Methods

- Notify neighboring land managers of invasive exotic plant occurrences.
- Offer to assist in exotic plant removal projects. Work with your local government to encourage the use of native plants in their urban and suburban landscapes.
- Provide them with lists of attractive, non-invasive, locally native alternatives that are naturally hardier, pest-resistant, and provide more nutritious food for wildlife than cultivated plants.

Hoeing or Hand Pulling

The hoeing or hand-pulling of small patches of weedy plants has been an effective method of controlling weeds since ancient times. It can prevent the spread of weeds to other areas if the seed heads and roots are bagged for proper disposal and not mulched into the soil.

Mowing

Cutting by hand, mowing with a mechanical mower, or whacking weeds with a string or blade trimmer before the weeds establish seeds may prevent the spread of the weeds to new areas. However, this method usually does not kill the weeds unless they are cut to the root line. This method still helps to control the spread of noxious weeds to new areas, but the mowing or cutting must be repeated periodically throughout the weed's growing period. Mowing must not be done when the weeds are bearing seed, unless the plants are caught in a basket or bag, or the seeds will be spread mechanically.

Solarization

Solarization of noxious weeds, over-shading or out-shading, is done with a thick layer of organic or inorganic mulch that may provide adequate shading and prevent growth of most light-sensitive plants. Most plants require sunlight for adequate growth. If heavy dark plastic is used, the edges must be secured with sandbags or rocks to hold the cover in place and to prevent sunlight from spreading in from the edges. This method may be very effective for controlling weeds in small patches. Solarization must be done at the beginning of the seed germination period. The solarizing cover must be left in place for the entire growing season for the noxious weed targeted. Solarization will kill both noxious weeds and native plants. Revegetation will be needed with fast-growing, hardy native seeds and plants to prevent the reintroduction of noxious weeds into the solarized areas and to prevent erosion of the bare soils left after solarization. Persistent follow-up and twice-yearly monitoring is needed to eliminate many of the invasive thistles, since some weedy thistle seeds can survive up to ten years.

Re-Seeding

Many weeds are opportunist that colonize disturbed areas where there is little competition from native plants. Re-seeding disturbed areas with competitive native plants is an effective long-term sustainable method of suppressing the weeds and allowing native plants to dominate weedy plants, choking out the weeds. Rapidly growing, competitive native plants should be used in weed control areas. Native

Noxious Weed Control Methods

plants do not require fertilizers or continued care, other than adequate water to establish their root systems. Native plants also provide forage for the local wildlife. Native plants should be seeded or planted at the beginning of the rainy season in the fall, usually around October 15th. Native plants recommended for re-seeding and replanting disturbed areas at the Tulloch Project are found in Tables at the end of this pamphlet. A list of potential native plant and native seed suppliers is also found at the end of this pamphlet.

Grazing

Livestock can readily feed on most seedling noxious weeds at Tulloch Reservoir. However, some of the noxious weeds at the Tulloch Project are toxic or harmful to livestock when mature, especially horses. Goats and sheep are more tolerant of these plants. Some of the seeds of the noxious weeds at Tulloch Reservoir can pass through the digestive tract of livestock, remain viable, and the seeds can be spread to new areas. Grazing livestock in close proximity to this drinking water reservoir can cause contamination with animal feces. Penning a flock of chickens, geese, or turkeys in small patches of seedy weeds for a few weeks can be an effective way of removing the entire year's seed production from that patch of seedy annuals. Seed-eating birds have gullets that can grind up seeds that would normally pass through livestock undigested. A drawback to using portable bird pens would be the potential predation of the birds by the fox and coyotes observed at Tulloch Reservoir. Livestock also eat desirable native plants. Livestock cannot be used in areas zoned by Calaveras or Tuolumne Counties for urban residential or commercial land uses, since livestock grazing is not permitted in these County zoning districts.

Burning

Controlled burns of weedy fields and spot burning of weeds with a propane torch have been reported successful for controlling many noxious weeds in other areas, but burning is not recommended at Tulloch Reservoir due to the following: the narrow band above the waters edge within the project area (below the 515-foot elevation); the project's proximity to residential and commercial structures; difficult access to the shoreline by fire suppression equipment; and the steep canyon surrounding the project area, where fire management would be extremely difficult. Contact your local fire department for burning regulations.

Noxious Weed Control Methods

Herbicides

For situations where other control methods of weeds are not practical or safe, herbicidal control has been reported as an effective option. Herbicides are likely also to damage desirable native plants and may harm animals. Applications can be made during the early spring when most other non-target vegetation is dormant. Herbicides are not appropriate for some infested sites, especially those near water, near Endangered plant habitats, or near Endangered wildlife habitats. Near streams or lakes, particular caution should be taken when using herbicides to avoid polluting the water. Remember that Tulloch Reservoir is used as a drinking water supply for the adjacent community.

Prior to using any herbicide, check with the County Agricultural Commissioner to determine which chemicals are legal to use in this situation. The labels also give more precise information on mixing and safety precautions. Refer to the pesticide manufacturer's label for specific information and restrictions regarding herbicide use. Read the entire herbicide label carefully. All label directions for mixing and applying must be followed. All precautions must be followed, and recommended personal protective clothing and gear must be used. Herbicide use should be done in consultation with a State-licensed Pest Control Advisor (PCA), registered to do such herbicide work in each County. Lists of such licensed PCA specialists are available from your County Agricultural Commissioner.

All pesticides registered for use have a label that has been approved by the Environmental Protection Agency (EPA). The label contains information about the product, including its relative toxicity, potential hazard to humans and the environment, directions for use, storage and disposal, and first aid treatment in case of exposure.

Contact the State Department of Pesticide Regulation for questions regarding the use requirements, restrictions, or recommendations for any herbicide to be used. <https://www.cdpr.ca.gov/>

Tulloch Reservoir Noxious Weeds

Barbed Goat Grass (*Aegilops triuncialis*)



Weed Description

Barbed Goatgrass is a winter annual closely related to and resembling winter wheat (*Triticum aestivum* L.). Stems are erect, spreading, or abruptly bent. Stems branch and spread at the base. Leaf blades are flat, spreading, about 0.1 inch wide, and may be sparsely covered with fine hairs. It grows to 18 inches (45 cm) tall. Stems are solid when young, but become hollow with age. Leaf blades are rigid and not susceptible to rust. Roots are fibrous and develop rapidly to a great depth. The flowering spikelets are 1 per node, are alternate and lay flat against a groove in the zig-zag tip of the stem. Immature spikes are often reddish or purplish. It has 2-3 stiff, sharp, and minutely barbed awns on each spikelet, each 1.5-3 inches long. There are two 1-seeded fruits per joint, with the upper two spikelets sterile. Spikelets adhere to the stem and are difficult to separate from the joint, falling at maturity with the entire flowering head.

Reproduction

The florets of Barbed Goatgrass bloom from May to July. They produce seeds. Barbed Goatgrass seeds germinate in the joints in the field, but joints appear to reduce germination. Germination occurs under a wide range of temperatures, 41-77 degrees F (5°-25° C). Some seeds can remain dormant for up to 5 years. The long barbed awns may attach and be transported on hair, fur, wool, shoes, or clothes.

Control

Some Barbed Goatgrass seeds falling on the ground can survive controlled field burns, because outer portions of the joints protect the seeds. Heavy grazing and high intensity/short duration grazing appears to increase goatgrass density. The awns can pierce and damage the mouth and eyes of livestock. Grazing animals should be removed from infested areas before plants mature. Mowing can be an effective method of reducing seed production. Mowing should occur after flowering, but before goatgrass seeds mature. Early mowing will result in new tiller growth and late mowing will only spread viable seed. Deep tilling has been effective. Hand pulling or hoeing small infestations is effective, if the roots are pulled and air-dried. Harvest transport trucks should be covered and other vehicles thoroughly cleaned before moving from infested fields. Nonselective herbicides such as Glyphosate may be applied to control small infestations. A reseeding program of perennial grasses and legumes, or other desirable vegetation should follow spot treatments.

Tulloch Reservoir Noxious Weeds

Chinese Tree-Of-Heaven (Ailanthus Altissima)



Weed Description

Chinese Tree-of-heaven, also known as *Ailanthus*, is a rapidly growing, deciduous tree native to Central China. *Ailanthus* has smooth stems with pale gray bark and chestnut brown twigs. It has large compound leaves, which range from 1-4 feet in length. These leaves are compound of 11-25 smaller leaflets, which are arranged alternately along the stems. Each leaflet has 1-3 coarse teeth and odor glands near the base. Mature trees can reach 80 feet or more in height. In the spring, small yellow-green flowers are found at the tips of branches. The seeds are enveloped by papery sheaths called samaras, which are slightly twisted, making them spin as they fall. Ripe samaras are greenish yellow or reddish brown in color. The wood of *ailanthus* is soft, coarse-grained, and creamy white to light brown in color. All parts of the tree, particularly the flowers, have a strong unpleasant odor, suggestive of rancid or burnt nuts.

Reproduction

Ailanthus reproduces both sexually via seeds and asexually via vegetative sprouts from the roots. Flowering occurs in June. *Ailanthus* is dioecious, with male and female flowering on

separate trees. Female trees produce seeds from late summer to early fall. A single tree can produce around 325,000 to 350,000 seeds each year. Seeds are found inside papery, twisted, winged structures called samaras, which are well adapted for wind dispersal. Samaras occur in large clusters from September to October, and may remain on the female tree through the winter, but can be dispersed any time from October to the following spring. Mature trees also produce suckers from the roots and re-sprout rapidly from cut stumps and root fragments.

Control

Due to its rapid growth, abundant seed production, high seed germination rate, and root and stump sprouting capabilities, the removal of *Ailanthus* requires more than one year of treatment and follow-up monitoring. Seedlings can be pulled or dug up, but removal or poisoning of the entire root system is important to prevent fragments from regrowing. Cutting *Ailanthus* trees may be ineffective unless the roots are also treated or poisoned, because the plant can produce stump sprouts and root suckers (connected to a pre-existing lateral root). Establishing a thick cover of native trees or shrubs will help shade out establishments of *Ailanthus* seedlings. The use of the herbicides Glyphosate, Triclopyr, Imazapyr, and Dicamba has been reported to be effective in *Ailanthus* control. Herbicides may be applied according to the state-approved pesticide labels to the leaves, the basal bark, cut stump, or the interior of the tree by injection.

Tulloch Reservoir Noxious Weeds

Italian Thistle (*Carduus pycnocephalus*)



Weed Description

The Italian Thistle is an annual, broadleaf weed with slender spiny-winged stems. It is typically 1-4 feet in height, but can grow up to 6 feet tall. The leaves are deeply cut into 5 pairs of spiny lobes. The upper surface of the leaf is smooth and the underside is covered with woolly hairs. Flower heads are covered with dense matted hairs. Its cylindrical flowers are purplish or pinkish in color. Flowers appear at the ends of branches, singly or in clusters of 2-5, and are about 12-3/4 inch long. Italian thistle has a branched, slender taproot.

Reproduction

Italian Thistle is bisexual and pollinated by many different insects. It reproduces only by two types of seeds - brown seeds and silver seeds. A single plant can produce 20,000 seeds. Brown seeds generally remain in the flower heads, falling with them to the ground at the end of the season, and germinate at lower temperatures than the silver seeds. Silver seeds have a large feathery pappus and are dispersed by wind. They can remain dormant in the soil longer than the brown seeds (8-10 years). Seeds can

germinate from depths of three inches, but usually germinate at less than one inch. Germination occurs in the fall after the first rain. Seedlings grow through the winter as rosettes and flower from May to July, before the summer drought. In addition to wind dispersal, seeds are also dispersed by contaminated hay or feed, water, animals, and machinery. Seedlings establish best on bare, disturbed soil.

Control

Mechanical removal such as digging will kill Italian Thistle, but the root must be cut at least four inches below ground level so the plant does not regrow. Plants can regrow from the base even if the thistle is repeatedly mowed at 3 inches. Plants should be pulled before seeds set. Grazing by sheep, goats, and horses has also been known to be effective in controlling this thistle. Three species of insects, *Psylliodes chalsomera*, *Rhinocyllus conicus*, and *Ceutorhynchus trimaculatus* damage vital parts of the thistle, impacting seed production. Concern that these insects may prey on several of California's endangered native thistles in the genus *Cirsium*. Infection by two fungal rusts, *Puccinia* and *Alternaria*, is known to significantly reduce growth of this Thistle. A variety of herbicides including Diquat, Picloram, Glyphosate, Clopyralid, Milestone and 2,4-D have been reported successful in the control of this plant.

Tulloch Reservoir Noxious Weeds

Slender-Flower Thistle (*Carduus tenuiflorus*)



Weed Description

Slender-flower Thistle is a broadleaf, winter annual weed with slender stems and is typically 1-6.5 feet in height. It has triangular shaped large stem wings with spiny tips. The leaves of this thistle are cut into 6-8 pairs of spine-tipped lobes and the undersides are covered with woolly hairs. The flowers are purplish or pinkish in color and appear at the ends of branches, typically in clusters of 5-20. This thistle resembles Italian Thistle, but Slender-flower thistle usually has more than five cylindrical flower heads per cluster, its bracts are not hairy, and it has leaves with 12-20 lobes. Seeds are gray-brown. Some references felt this thistle is an indistinct species from Italian Thistle.

Reproduction

Slender-flower Thistles reproduce only by seed. Germination occurs in the fall after the first rain when temperatures reach 36-86 degrees F (2-30 degrees C [Centigrade]). Seedlings grow through the winter as rosettes, and flower from May-July, before the summer drought. Its small seeds have

a large pappus and are well adapted for wind dispersal. Seeds are also dispersed by contaminated hay or feed, water, animals, and machinery.

Control

Mechanical removal such as digging will kill Slender-flower Thistle, but the root must be cut at least four inches below ground level so the plant does not regrow. Plants can regrow from the base even if the thistle is repeatedly mowed at 3 inches. Plants should be pulled before seeds set. Grazing by sheep, goats, and horses has also been known to be effective in controlling this thistle. Three species of insects, *Psylliodes chalsomera*, *Rhinocyllus conicus*, and *Ceutorhynchus trimaculatus* damage vital parts of the thistle, impacting seed production. Infection by two fungal rusts, *Puccinia* and *Alternaria*, is known to significantly reduce growth of this Thistle. A variety of herbicides including Diquat, Picloram, Glyphosate, Clopyralid, Milestone and 2,4-D have been reported successful in the control of this plant.

Tulloch Reservoir Noxious Weeds

Tocalote (Malta Starthistle) (*Centaurea melitensis*)



Weed Description

Tocalote is an annual or biennial thistle with several stems from the base and branched in the upper part. Its erect or drooping stems have narrow wings and are 4-36 inches high. Most leaves are thin oblongs with deep lobes, 2-4 inches long. Leaves are alternate on the stem. Flowers are yellow and 0.3-0.5 inches long. Flower heads are in the upper leaf axils and are globe-shaped with firm bracts that are hairy when young. The margins of the bracts have several very short tan spines near the tip, with reddish, pinnately spiny bases. The seed pappus bristles are in several series. The similar weed, Yellow Star Thistle, has longer, yellow spines.

Reproduction

Tocalote blooms from May to June. Tocalote is insect-pollinated and reproduces by seed. Seed production is highly variable. Plants can produce 1-60 or more seeds per head and 1-100 or more heads per plant. Seeds fall near the parent plant or are dispersed to short distances by the wind, water, and soil movement. Seeds are often

Spread by humans, machinery, and livestock via the transportation of contaminated crop seed or hay. Seeds are also spread by road maintenance and moving livestock, which can contribute to the long-distance spread of the seeds. Wild birds such as quail and finches feed heavily on the seeds and are capable of long-distance dispersal. The seed bristles have microscopic barbs that can adhere to clothing, hair, and fur for seed dispersal. Seeds germinate after the first fall rains. Plants exist as basal rosettes through winter and early spring until flowering stems develop. Young seedlings are especially resistant to the effects of fall drought.

Control

Methods used to control Yellow Star Thistle can be applied to Tocalote. High-intensity, short-duration grazing by goats or sheep can be implemented. Mowing and spot burning can be very effective if done before spines form on the flower heads. Mowing too early can result in higher seed production. Monitoring and spot eradication of plants when they are discovered can also prevent the spread of this thistle. Hand pulling or grubbing, removing as much root as possible, may be useful in eradicating small infestations. Re-seeding with competitive native plant species is an effective long-term sustainable method of outcompeting Tocalote. Wild oat (*Avena* spp.) litter appears to inhibit Tocalote germination. The herbicides Clopyralid, Milestone, and Picloram have been reported to be effective in controlling Tocalote.

Tulloch Reservoir Noxious Weeds

Yellow Star Thistle (*Centaurea solstitialis*)



Weed Description

Yellow Star Thistle is gray-green to blue-green in color. It grows from 0.5 to 3 feet in height and has deep taproots. Flowers are bright yellow, with long, sharp spines surrounding the flower base. Stems and leaves are covered with soft wool. Basal leaves are deeply lobed and 2-3 inches in length. Upper leaves are sharply pointed and only 0.5-1.0 inches in length.

Reproduction

Yellow Star Thistle is a non-woody annual plant. It flowers from late May to September, sometimes into December or later. With low levels of self-fertilization, this thistle depends on honeybees, bumblebees, and other insects for pollination. Each seed head can produce approximately 35 to 80 seeds. Pappus-bearing seeds are dispersed after the flowers drop their petals, while non-pappus-bearing seeds may remain on the plant well into the winter. Without effective wind dispersal mechanisms, most seeds fall to the ground at the base of the parent plant. Even the plant's pappus-bearing seeds fall within 2 feet of the parent plant 92% of the time.

Dispersal over greater distances is assisted by animals and human beings. Birds such as house finches, goldfinches, pheasants, and quail feed on its seeds and transport them far from the parent plant. Long-distance dispersal of the seeds is also assisted by vehicles, contaminated hay or crop seed, and moving livestock. Germination occurs throughout the rainy season (October - June), with the maximum germination after the early fall rains when the seeds are exposed to moisture, light, and favorable temperatures of 50-68 F (10-20°C). Seeds remain dormant until they have adequate amounts of moisture.

Control

For small patches, manual removal works well. Hand-pulling or weed-whacking is best done before full flowering or viable seed production (May-June). Grazing by cattle, sheep, or goats can reduce abundance, growth, and productivity when done during May-June, after the plants have bolted, but before the buds produce spines. Toxic to horses. Re-seeding with competitive native plant species is an effective long-term sustainable method of outcompeting *C. solstitialis*. Competitive grasses used in re-vegetation programs for thistle management include native perennial grasses, such as big bluegrass (*Poa ampla*) and thick-spike wheatgrass (*Elymus lanceolatus*). The herbicides Clopyralid, Picloram, 2,4-D, Dicamba, Milestone, and Triclopyr have been reported to be effective in controlling Yellow Star Thistle as well.

Tulloch Reservoir Noxious Weeds

Bull Thistle (*Cirsium vulgare*)



Weed Description

The Bull Thistle is a biennial plant 2-5 feet in height. Stems are erect, hairy, spiny-winged, and branched. Lance-shaped leaves are short and broad, usually measuring less than 12 inches in length, and are very wavy or crinkled. They are prickly above but cottony underneath, giving the underside a grayish appearance. The leaves are lobed with 3-4 points per lance-shaped lobe, and each point ends in a long yellow spine with shorter spines between. In the first year, leaves form a rosette. Mature leaves alternate along the stem, causing the stalk to be prickly and winged. Flowers are found at the tips of branches, singly or in clusters. They are pinkish-purple and 1 ½ - 2 inches wide. Flower-head bracts are covered with spines. The pappus-bearing seeds of the Bull Thistle are oblong and straw colored.

Reproduction

The Bull Thistle flowers from June to September. It reproduces solely by seed. Mature plants can produce up to 4,000 seeds each. Seeds are typically 5 mm long and have a downy pappus

making them well suited for wind dispersal. Establishment is promoted by soil disturbance, which increases nutrient, water, and light availability to seedlings and reduces the vigor of competing vegetation

Control

The best control measure for Bull Thistle is to prevent the production and movement of seeds. Cleaning equipment before leaving infested areas, feeding hay that is not contaminated, and planting weed-free seed will prevent the spread of this thistle. Mowing or hand-cutting before the production of viable seeds will also prevent the spread of Bull Thistle. Mowing should be done after the pappus of the earliest seeds displays color and before they turn brown. When digging, the root should be dug at least 2 inches below ground level. Heads that are beyond the bud stage should be selected and placed in a tight container to be buried in a landfill. The Government has approved a biological control for Bull Thistle; however, it may attack other native *Cirsium* thistles. The herbicides 2,4-D Amine, Chlorsulfuron, Picloram, and Dicamba have been reported effective in the control of Bull Thistle when applied during the rosette stage of growth. Other herbicides reported successful in the control of this plant are Metsulfuron, Imazapic, Clopyralid, Milestone, and Triclopyr.

Tulloch Reservoir Noxious Weeds

Klamath Weed (St. John's Wort) (*Hypericum perforatum*)



September, with peak flowering occurring from mid to late June. Plants typically produce an average of 15,000-30,000 seeds each season. Seeds may be produced without fertilization. Developing seed capsules become moist and sticky. These sticky capsules can disperse by wind or water or by adhering to animals and humans. Seeds are hard-coated and remain intact even after being consumed by seed predators. Seeds may remain viable in the soil for up to 10 years. Germination occurs from fall through spring. Brief exposure to fire (212-284° F) often increases germination, while small amounts of calcium may affect the permeability of the seed coat and inhibit germination.

Weed Description

Klamath Weed stands 1 to 4 feet tall and has numerous rust-colored branches. Leaves are green, opposite, oblong-shaped, covered with tiny transparent dots, and are usually no more than an inch in length. Flowers have five separate petals and are bright yellow in color. The petals are about 0.5 inches long and dotted with black glands around the edges of each petal. Yellow colored stamens are arranged in three groups. The seeds are shiny black to brown and approximately 1 mm long. Seeds are often coated with a sticky, gel-like material that assists in dispersal. The taproot of this perennial plant may reach depths of 4 to 5 feet and allows it to survive year-round. Lateral roots typically grow 2 to 3 inches beneath the soil surface.

Reproduction

Klamath Weed is a perennial plant that reproduces by seed and by vegetative sprouts. Root systems are capable of forming buds, which grow into new plants separating from the parent plant. Klamath Weed flowers from June to

Control

Klamath Weed is known to be poisonous to animals, including causing brain lesions in livestock, and is generally avoided by grazers. So, grazing is not recommended for control. However, several beetles, including *Chrysolina hyperici* and *Chrysolina quadragemina*, feed on the foliage and have been used successfully as biological controls. Mechanical removal, such as mowing of Klamath weed, may stimulate vegetative reproduction from the roots. Herbicides that have been reported as effective for control of this *Hypericum* include 2,4-D, Picloram, and Glyphosate.

Tulloch Reservoir Noxious Weeds

Medusa Head (*Taeniantherum* [*Elymus* caput-medusae])



Weed Description

Medusa-head is an aggressive winter annual grass with slender stems. It reaches 6-24 inches in height. Leaf blades are approximately 1-3 mm (<0.1 inches) wide and roll inward at the margins. There are typically 24 blades per stem. The awns or spikes at the tips of the plant are 1-4 inches long, minutely barbed, are compressed into the center when green, and twist apart upon drying. This species matures 2-4 weeks later than most other annual grasses and stands out green against surrounding brown grasses. After producing seeds, Medusa-head turns light gray or tan in color. The seed heads do not break apart as seeds mature.

Reproduction

Medusa-head reproduces by seed. Seeds are dispersed by wind, water, and also by attaching to the coats of animals, machinery, vehicles, and clothing. Medusa-Head flowers in May-June and is predominantly self pollinating. An average of 8-15 seeds are produced per spike in late spring or early summer. Seeds usually disperse by mid-summer and can remain dormant for over six months. Seeds typically germinate after the first

rains in October or November, and the germination rate is high. Germination occurs under a broad temperature range (optimal 10-150 C or 50-59 F). The seeds of this plant are capable of germinating and growing in the dense leaf litter layer, without making contact with the soil. If they suffer desiccation, the primary root dies, but a new root develops if they are moistened again. Seeds can emerge from soil depths of up to 3-4 inches. Medusa-head continues to grow and produce seeds later than most grasses.

Control

Mowing for two consecutive years can help reduce infestations of Medusa-head. Mowing in combination with grazing, was found to be effective in reducing infestations. Heavy grazing by sheep in early spring (when it is still palatable) can assist in controlling it, but animals should be removed before seed heads form to limit dispersal. The stiff awns and hard florets can injure eyes and mouths of grazing animals. Plowing or disking before seeds set are also effective means for control. Revegetation with competitive native species, like Squirreltail grass (*Elymus elymoides*), may prevent Medusa-head from regaining dominance after control treatments. Burning stands of Medusa-head prior to seed dispersal is reported as an effective control measure. The western U.S. endemic crown rot fungus (*Fusarium culmorum*) is a potential biocontrol agent adapted to dry soils. The herbicides Atrazine, Glyphosate, and Paraquat have been reported effective to varying degrees, though they may cause harm to other species.

References

Balls, Edward K. , "Early Uses of California Plants", University of California Press, 1962.

Barr, C.B.. U.S. Department of the Interior. Fish and Wildlife Service, Sacramento, CA; "The distribution, habitat, and status of the Valley Elderberry Longhorn Beetle *Desmocerus californicus dimorphus*", 1991.

Blackwell, Laird R.. Wildflowers Of the Sierra Nevada and the Central Valley', 1999.

Brown, Georgia H. , "Plants Used by Indians and Early Californians for Food and Medicine", manuscript at San Luis Obispo County Schools, 1948.

Burke, HE. , "Biological Notes on *Desmocerus*, a Genus of Roundhead borers, the Species Of which Infest Various Elderberries", Journal of Economic Entomology, 14450-452, 1921.

Brown, Lauren, "Grasses, An Identification Guide", Roger Tory Peterson Institute, 1979.

California Department of Food & Agriculture, Division of Plant Health & Pest Prevention Services, "Biological Control Program Annual Report, 2003.

California Department of Food and Agriculture, Encyclopedia of Weeds. April 2006.

California Department of Food & Agriculture, Division of Plant Health and Pest Prevention Services, "Pest Ratings of Noxious Weed Species and Noxious Weed Seecf, February 2005.

California Department of Pesticide Regulation, "What You Need to Know About Protecting Rivers from Dormant Spray Residues", March 2003.

California Department of Pesticide Regulation, "Pesticides by Species (Volume II), An Index to Pesticides That Are Used in Proximity to Federally Listed, Proposed and Candidate Species in California By Active Ingredient", August 1997

California Invasive Plant Council (CAL IPC), "California Invasive Plant Inventory", February 2006.

California Invasive Plant Council (CAL IPC), •Exotic Pest Plants of Greatest Ecological Concern in California" list, 1999.

California Native Plant Society, "A Manual of California Vegetation"; J. O. Sawyer and T. Keeler-Wolf, 1995.

Clarke, Charlotte B., "Edible and Useful Plants of California", California Natural History Guides, 1977.

Crampton, Beecher, "Grasses in California", 1974.

"Federal Noxious Weed Regulation", List 3, 7 CFR 360, 2004.

Ferris, Roxana S.. "Native Shrubs Of the San Francisco Bay Region", 1968.

Fuller, Thomas C. & Elizabeth McClintock, "Poisonous Plants of California", University of California press, 1986.

Hickman, J. C, Editor, "The Jepson Manual, Higher Plants Of California", University Of California Press, 1993.

References

Holland. R.F., "Preliminary Description of the Terrestrial Natural Communities of California", Unpublished Report of the State Department of Fish and Game, Natural Heritage Division, Sacramento, CA, 1986.

Lanzara, Paola and Mariella Pizzetti, "Simon & Schuster's Guide to Tæes", edited by Stanley Schuler, 1978.

Munz, Philip A.. "A California Flora", University of California Press, 1970.

Natural Diversity Database, Division Of the State Department of Fish and Game, 2006.

Niehaus, Theodore and Charles Ripper, "The Peterson Field Guide series, A Field Guide to Pacific States Wildflowers. National Audubon Society and National Wildlife Federation, 1976.

Pahlow, Mannfried, "Healing Plants." Barron's Educational Series, Inc., 1993.

Schoenig, Editor Steve, "California Noxious & Invasive Weed Action Plan, California Department of Food & Agriculture and California Invasive Weed Awareness Coalition, September 2005.

Spellenberg, Richard, keys by Susan Rayfield and Carol Nehring, The Audubon Society Field Guide to North American Wildflowers, Western Region, 1979.

Tibor, David, Editor, •California Native Plant Society Inventory of Rare and Endangered Plants of California", California Native Plant Society, 2001.

Tri-Dam Project, Tulloch Project FERC Project No. 2067, Volume II Application for New License for the relicensing of the Tulloch Hydroelectric Project, 2002.

Tri-Dam Project, "Final Environmental Impact Statement Stanislaus River Projects" for the relicensing of the Tulloch Hydroelectric Project, March 2005.

Tu, Mandy, Callie Hurd, & John M. Randall, "Weed Control Methods Handbook: Tools and Techniques for Use in Natural Areas", published by The Nature Conservancy and available at the internet <http://trmweeds.ucdavis.edu/handbook>, April 2001.

University of California Agricultural and Natural Resources Department, •Yellow Star-thistle, Integrated Pest Management of Land Managers, Landscape Professionals, and Home Gardeners", Pest Notes, Publication 7402. July 2003.

U.S. Environmental Protection Agency, "Protecting Endangered Species Interim Measures for Use of Pesticides in Calaveras County", March 2000.

U.S. Environmental Protection Agency, "Protecting Endangered Species Interim Measures for Use of Pesticides in Tuolumne County", March 2000.

U. S. Department of the Interior, Fish and Wildlife Service, Endangered Species Program, Portland, Oregon, "Recovery Plan for the Valley Elderberry Longhorn Beetle", 1984.